arXiv:1607.05055v1 [math.CV] 18 Jul 2016

FEKETE POLYNOMIALS AND SHAPES OF JULIA SETS

KATHRYN A. LINDSEY AND MALIK YOUNSI

Abstract. We prove that a nonempty, proper subset S of the complex plane can be approximated in a strong sense by filled Julia sets of polynomials of degree at least two if and only if S is bounded and $\hat{\mathbb{C}} \setminus \text{int}(S)$ is connected. The proof that such a set is approximable by filled Julia sets is constructive and relies on Fekete polynomials. Illustrative examples are presented. We also prove an estimate for the rate of approximation in terms of geometric and potential theoretic quantities.

1. Introduction

The study of possible shapes of polynomial Julia sets was instigated by the first author in [10], who proved that any Jordan curve in the complex plane $\mathbb{C}$ can be approximated by polynomial Julia sets arbitrarily well in the Hausdorff distance, thereby answering a question of W.P. Thurston.

Theorem 1.1 (Lindsey [10])**.** *Let $E \subset \mathbb{C}$ be any closed Jordan domain. Then for any $\epsilon > 0$, there exists a polynomial P such that*

$$d(E, \mathcal{K}(P)) < \epsilon, \quad d(\partial E, \mathcal{J}(P)) < \epsilon.$$

Here $\mathcal{K}(P) := \{z \in \mathbb{C} : P^m(z) \not\to \infty \text{ as } m \to \infty\}$ is the filled Julia set of P, $\mathcal{J}(P) := \partial \mathcal{K}(P)$ is the Julia set of P and d is the Hausdorff distance.

Note that Theorem 1.1 remains valid if E is any nonempty connected compact set with connected complement, by a simple approximation process.

Approximation of planar sets by Julia sets has subsequently been the focus of several other works, such as [2], where it was shown that any connected compact set in the plane can be approximated by dendrite Julia sets. See also [9] for applications to computer graphics.

An important feature of the approach in [10] which seems to be absent from other works is that it is constructive and can easily be implemented to obtain explicit images of Julia sets representing various shapes.

In this paper, we prove the following generalization of Theorem 1.1.

Theorem 1.2. *Let $E \subset \mathbb{C}$ be any nonempty compact set with connected complement. Then for any $\epsilon > 0$, there exists a polynomial P such that*

$$d(E, \mathcal{K}(P)) < \epsilon, \quad d(\partial E, \mathcal{J}(P)) < \epsilon.$$

Note that here E is not assumed to be connected.

Date: July 14 2016.

2010 *Mathematics Subject Classification.* primary 30E10, 37F10; secondary 30C85.

Key words and phrases. Fekete points, polynomials, Julia sets, Hausdorff distance, Leja points.

Second author supported by NSERC.

This supersedes [10, Theorem 4.3], which deals with rational maps instead of polynomials.

The proof of Theorem 1.2 generalizes the method introduced in [10] by putting it in the more natural framework of potential theory. More precisely, our approach is based on the observation that in Theorem 1.1, the roots of the polynomial P are equidistributed with respect to harmonic measure on ∂E. In the connected case, such points can be obtained as images of equally spaced points on the unit circle under the Riemann map. This, however, does not extend to disconnected sets and in this case, the explicit construction of equidistributed sequences of points is more difficult. Nevertheless, this is a classical problem in potential theory which has been extensively studied in the past. For instance, Fekete described in [5] the following construction :

Let E be a nonempty compact set with connected complement, and let $n \geq 2$. A *Fekete n-tuple* for E is any n-tuple $w_1^n, \ldots, w_n^n \in E$ which maximizes the product

$$\prod_{j<k} |w_j^n - w_k^n|.$$

Note that by the maximum principle, any Fekete n-tuple lies in ∂E.

Fekete points are classical objects of potential theory which have proven over the years to be of fundamental importance to a variety of problems related to polynomial interpolation. For instance, they can be used to give a proof of Hilbert's Lemniscate Theorem, which states that any compact plane set with connected complement can be approximated by polynomial lemniscates. In fact, as we will see, Hilbert's theorem is closely related to Theorem 1.2. For more information on Fekete points and their applications, we refer the reader to [4] and [16, Chapter III, Section 1].

It is not difficult to show that Fekete points are indeed equidistributed with respect to harmonic measure on ∂E, in the sense that the counting measures

$$\mu_n := \frac{1}{n} \sum_{j=1}^{n} \delta_{w_j^n}$$

converge weak* to the harmonic measure (see Proposition 2.12). Combining a stronger version of this result due to Pritsker [14] with the method put forward in [10] gives a constructive proof of Theorem 1.2.

Other equidistributed sequences of points include Leja points, which share many nice properties with Fekete points but are easier to compute numerically. Theorem 6.3 proves that our approximation technique works if Leja points, instead of Fekete points, are used. The technique in [10] for constructing polynomials whose filled Julia sets approximate a given Jordan domain required first obtaining an approximation of the Riemann map on the complement of that domain. Using Leja points or Fekete points obviates this step, and thus constitutes an advance in the constructive process.

Note that Theorem 1.2 not only yields approximation of the set E by the filled Julia set of a polynomial, but also approximation of its boundary by the corresponding Julia set. Consequently, we introduce the following definition.

Definition 1.3. A nonempty proper subset S of the plane is *totally approximable* by a collection $\mathcal{C}$ of nonempty proper subsets of the plane if for any $\epsilon > 0$, there

exists $C \in \mathcal{C}$ such that

$$d(S, C) < \epsilon, \quad d(\partial S, \partial C) < \epsilon.$$

Note that closeness of two sets in the Hausdorff distance neither implies nor is implied by closeness of their boundaries, see Figure 1. On the other hand, it is easy to see that closeness of both the sets and their boundaries implies closeness of the complements, c.f. Lemma 5.4.



FIGURE 1. A unit disk, a unit disk with a small disk removed, and a ring of small disks. The first two are close in the Hausdorff distance, but their boundaries are not. The boundaries of the first and third are close in the Hausdorff distance, but the sets themselves are not.

With this definition, Theorem 1.2 states that any nonempty compact plane set E with connected complement is totally approximable by polynomial filled Julia sets. More generally, using an approximation process, one can prove that this remains true if E is replaced by any nonempty bounded set whose interior has connected complement. This condition also turns out to be necessary.

Theorem 1.4. *A non-empty proper subset S of the complex plane $\mathbb{C}$ is totally approximable by filled Julia sets of polynomials of degree at least 2 if and only if S is bounded and $\mathbb{C} \setminus \operatorname{int}(S)$ is connected.*

Finally, our method also gives a precise estimate for the rate of approximation in Theorem 1.2. More precisely, let $E \subset \mathbb{C}$ be a compact set with connected complement Ω in the Riemann sphere $\widehat{\mathbb{C}}$. We also assume that the interior of E is not empty. In particular, the Green's function $g_\Omega(\cdot, \infty)$ for Ω with pole at ∞ exists. For $n \in \mathbb{N}$, define $s_n(E)$ to be the infimum of $s > 0$ for which there exists a polynomial p_n of degree n such that

$$E \subset \mathcal{K}(p_n) \subset E_s,$$

where $E_s := E \cup \{z \in \Omega : g_\Omega(z, \infty) \leq s\}$.

Theorem 1.5. *Let $E \subset \mathbb{C}$ be a uniformly perfect compact set with nonempty interior and connected complement. Then there exists a real number $c = c(E)$ depending only on E such that*

$$s_n(E) \leq c \, \frac{\log n}{\sqrt{n}} \qquad (n \geq 5).$$

Recall that a compact subset E of the plane is *uniformly perfect* if there exists a real number $a > 0$ such that for any $z \in E$ and for any $0 < r < \operatorname{diam}(E)$, there

is a point $w \in E$ with $ar \leq |z - w| \leq r$. This condition on E is not too restrictive, since, for example, uniformly perfect sets include compact sets consisting of finitely many nontrivial connected components.

A direct consequence of Theorem 1.5 is that $s_n(E) \to 0$ as $n \to \infty$. In particular, Theorem 1.5 implies Theorem 1.2 for uniformly perfect sets, since the compact sets E_s shrink to E as s decreases to 0. We also mention that results similar to Theorem 1.5 were obtained in [1] and [7], but for the rate of approximation by polynomial lemniscates in Hilbert's Lemniscate Theorem.

The remainder of the paper is organized as follows. Section 2 contains various preliminaries from potential theory, including the properties of Fekete polynomials. In Section 3, we prove an explicit version of Theorem 1.2 which has Theorem 1.5 as a corollary. Then, in Section 4, we give an alternate proof of Theorem 1.2 using Hilbert's Lemniscate Theorem. Section 5 contains the proof of Theorem 1.4. Lastly, Section 6 is devoted to numerical examples. We first discuss Leja points and prove that our approximation scheme remains valid if Fekete points are replaced by Leja points. This allows us to compute images of Julia sets representing various disconnected shapes.

2. Preliminaries from potential theory

This section contains various preliminaries from potential theory, including the notions of potential, logarithmic capacity, Green's function and Fekete points. The material is quite standard and is treated thoroughly in [15] and [16] for example, but we include it for the reader's convenience.

We begin with the notions of potential and energy of a measure.

Definition 2.1. Let μ be a finite positive Borel measure on $\mathbb{C}$ with compact support.

(i) The *potential* of μ is the function $p_\mu : \mathbb{C} \to [-\infty, \infty)$ defined by

$$p_\mu(z) := \int \log|z - w| \, d\mu(w) \qquad (z \in \mathbb{C}).$$

(ii) The *energy* of μ, noted $I(\mu)$, is given by

$$I(\mu) := \int \int \log|z - w| \, d\mu(z) \, d\mu(w) = \int p_\mu(z) \, d\mu(z).$$

One can check that p_μ is subharmonic on $\mathbb{C}$, harmonic on $\mathbb{C} \setminus (\operatorname{supp} \mu)$ and that

$$p_\mu(z) = \mu(\mathbb{C}) \log|z| + O(|z|^{-1}) \tag{1}$$

as $z \to \infty$.

Now, let E be a nonempty compact subset of $\mathbb{C}$. We assume that the complement of E is connected. Let $\mathcal{P}(E)$ denote the collection of all Borel probability measures supported on E.

Definition 2.2. A measure $\nu \in \mathcal{P}(E)$ is an *equilibrium measure* for E if

$$I(\nu) = \sup_{\mu \in \mathcal{P}(E)} I(\mu).$$

A standard weak*-convergence argument shows that E always has an equilibrium measure, see [15, Theorem 3.3.2]. Moreover, if the supremum in the definition is not $-\infty$, then the equilibrium measure ν for E is always unique and supported on

∂E ([15, Theorem 3.7.6]). It also coincides with $\omega_\Omega(\cdot, \infty)$, the harmonic measure for $\Omega := \widehat{\mathbb{C}} \setminus E$ and ∞.

The energy of the equilibrium measure is used to define the logarithmic capacity of E.

Definition 2.3. The *logarithmic capacity* of E is defined by

$$\mathrm{cap}(E) := e^{I(\nu)},$$

where ν is the equilibrium measure for E.

Remark. The quantity $-I(\nu) = -\log \mathrm{cap(E)}$ is usually called *Robin's constant*.

For example, the capacity of a closed disk is equal to its radius and the capacity of a segment is equal to a quarter of its length.

For the rest of the paper, we shall suppose that $\mathrm{cap}(E) > 0$. This can always be assumed without loss of generality in Theorem 1.2, adjoining a small disk in E if necessary. The condition of positive logarithmic capacity is enough to ensure that the Green's function exists.

Definition 2.4. Let $\Omega := \widehat{\mathbb{C}} \setminus E$ be the complement of E in the Riemann sphere. The *Green's function* for Ω with pole at ∞ is the unique function $g_\Omega(\cdot, \infty) : \Omega \to (0, \infty]$ such that

(i) $g_\Omega(\cdot, \infty)$ is harmonic on $\Omega \setminus \{\infty\}$;
(ii) $g_\Omega(\infty, \infty) = \infty$ and as $z \to \infty$,

$$g_\Omega(z, \infty) = \log|z| + O(1);$$

(iii) $g_\Omega(z, \infty) \to 0$ as $z \to \zeta$, for all $\zeta \in \partial\Omega$ except possibly a set of zero logarithmic capacity.

The Green's function for Ω with pole at ∞ can be recovered from the potential of the equilibrium measure ν by the following formula :

$$g_\Omega(z, \infty) = p_\nu(z) - I(\nu) \qquad (z \in \Omega \setminus \{\infty\}).$$

This is usually referred to as Frostman's theorem ([15, Theorem 3.3.4]). Combining this with Equation (1), we get

$$I(\nu) = \log \mathrm{cap}(E) = \lim_{z \to \infty} (\log|z| - g_\Omega(z, \infty)).$$

We will also need the notion of Harnack distance.

Definition 2.5. Let D be a domain in $\widehat{\mathbb{C}}$. Given $z, w \in D$, the *Harnack distance* between z and w is the smallest number $\tau_D(z, w)$ such that, for every positive harmonic function u on D,

$$\tau_D(z, w)^{-1} u(w) \leq u(z) \leq \tau_D(z, w) u(w).$$

The existence of τ_D is a simple consequence of Harnack's inequality. It is not difficult to show that $\log \tau_D(z, w)$ is a continuous semimetric on D ([15, Theorem 1.3.8]).

We now introduce a discrete approach to logarithmic capacity.

Definition 2.6. For $n \geq 2$, the *n-th diameter* of E is given by

$$\delta_n(E) := \sup_{w_1, \ldots, w_n \in E} \prod_{j<k} |w_j - w_k|^{\frac{2}{n(n-1)}}.$$

Recall from the introduction that an n-tuple $w_1^n, \dots, w_n^n \in E$ for which the supremum is attained is called a Fekete n-tuple for E.

Since E is compact and nonempty, a Fekete n-tuple always exists, though it need not be unique. Moreover, the maximum principle shows that any Fekete n-tuple lies in ∂E. Also, the quantity $\delta_2(E)$ is the usual diameter of E, and $\delta_n(E) \leq \delta_2(E)$ for all n. In fact, the sequence $(\delta_n(E))_{n\geq 2}$ is decreasing and has a limit, which is nothing other than the logarithmic capacity of E.

Theorem 2.7 (Fekete–Szegö)**.** *The sequence $(\delta_n(E))_{n\geq 2}$ is decreasing and*

$$\lim_{n\to\infty} \delta_n(E) = \operatorname{cap}(E).$$

See [15, Theorem 5.5.2].

The rest of this section is devoted to the properties of Fekete polynomials.

Definition 2.8. A *Fekete polynomial* for E of degree n is a polynomial of the form

$$q(z) := \prod_{j=1}^{n} (z - w_j^n),$$

where $w_1^n, \dots, w_n^n$ is a Fekete n-tuple for E.

In the following, we denote by $\|q\|_E$ the sup-norm of the polynomial q on E, i.e.

$$\|q\|_E := \sup_{z\in E} |q(z)|.$$

Theorem 2.9. *Let q be a monic polynomial of degree n. Then*

$$\|q\|_E^{1/n} \geq \operatorname{cap}(E).$$

If in addition q is a Fekete polynomial for E, then

$$\|q\|_E^{1/n} \leq \delta_n(E).$$

Proof. To prove the first statement, it suffices to observe that since $E \subset q^{-1}\left(\overline{\mathbb{D}}(0, \|q\|_E)\right)$, we have

$$\operatorname{cap}(E) \leq \operatorname{cap}\left(q^{-1}\left(\overline{\mathbb{D}}(0, \|q\|_E)\right)\right),$$

by the monotonicity of logarithmic capacity. The quantity on the right-hand side is easily seen to be equal to $\operatorname{cap}\left(\overline{\mathbb{D}}(0, \|q\|_E)\right)^{1/n} = \|q\|_E^{1/n}$, since in this case the Green's function can be identified explicitly. This proves the first statement.

To prove the second statement, write $q(z) := \prod_{j=1}^{n}(z - w_j^n)$ where $w_1^n, \dots, w_n^n$ is a Fekete n-tuple for E. If $z \in E$, then $z, w_1^n, \dots, w_n^n$ is an $(n+1)$-tuple in E, so

$$\prod_{i=1}^{n} |z - w_i^n| \prod_{j<k} |w_j^n - w_k^n| \leq \delta_{n+1}(E)^{n(n+1)/2},$$

and hence

$$|q(z)| \leq \frac{\delta_{n+1}(E)^{n(n+1)/2}}{\delta_n(E)^{n(n-1)/2}} \leq \frac{\delta_n(E)^{n(n+1)/2}}{\delta_n(E)^{n(n-1)/2}} = \delta_n(E)^n,$$

as required.

□

Theorem 2.10 (Bernstein's Lemma)**.** *Let E be as before, and let $\Omega = \widehat{\mathbb{C}} \setminus E$. If q is a polynomial of degree n, then*

$$\left(\frac{|q(z)|}{\|q\|_E}\right)^{1/n} \leq e^{g_\Omega(z,\infty)} \qquad (z \in \Omega).$$

If in addition q is a Fekete polynomial for E, then

$$\left(\frac{|q(z)|}{\|q\|_E}\right)^{1/n} \geq e^{g_\Omega(z,\infty)} \left(\frac{\operatorname{cap}(E)}{\delta_n(E)}\right)^{\tau_\Omega(z,\infty)} \qquad (z \in \Omega).$$

Proof. To prove the first statement, assume without loss of generality that q is monic, and define

$$u(z) := \frac{1}{n}\log|q(z)| - \frac{1}{n}\log\|q\|_E - g_\Omega(z,\infty) \qquad (z \in \Omega \setminus \{\infty\}).$$

Then u is subharmonic on $\Omega \setminus \{\infty\}$. Also,

$$u(z) = \log|z| - \frac{1}{n}\log\|q\|_E - \log|z| + \log\operatorname{cap}(E) + o(1)$$

as $z \to \infty$, and therefore setting

$$u(\infty) := \log\operatorname{cap}(E) - \frac{1}{n}\log\|q\|_E$$

makes u subharmonic on Ω. Now, since $\partial\Omega \subset E$ and since $u(\infty) \leq 0$ by Theorem 2.9, we have

$$\limsup_{z\to\zeta} u(z) \leq 0$$

for all $\zeta \in \partial\Omega$, and thus $u \leq 0$ on Ω by the maximum principle for subharmonic functions. This implies the result.

To prove the second statement, note that if q is a Fekete polynomial for E, then in particular all its zeros lie in E, so that u is actually harmonic on Ω. Also, from the first part of the proof, we have $u \leq 0$ on Ω. Therefore, applying the definition of Harnack distance to $-u$, we get

$$u(z) \geq \tau_\Omega(z,\infty)u(\infty) \qquad (z \in \Omega).$$

Now, by Theorem 2.9,

$$u(\infty) = \log\operatorname{cap}(E) - \frac{1}{n}\log\|q\|_E \geq \log\operatorname{cap}(E) - \log\delta_n(E).$$

Combining the last two inequalities yields the desired conclusion.

□

Corollary 2.11. *For $z \in \Omega$, we have*

$$|z| \leq R(E)e^{g_\Omega(z,\infty)},$$

where $R(E) := \sup_{w\in E}|w|$.

Proof. This follows from the first part of Theorem 2.10 applied to $q(z) := z^n$.

□

The following result shows that Fekete points are a good discrete approximation of the equilibrium measure.

Proposition 2.12. *Let E be as before and for each $n \geq 2$, let*

$$\mu_n := \frac{1}{n}\sum_{j=1}^{n} \delta_{w_j^n},$$

where $w_1^n, w_2^n, \ldots, w_n^n$ is a Fekete n-tuple for E and $\delta_{w_j^n}$ is the unit point mass at the point w_j^n. Then $\mu_n \to \nu$ weak*, where ν is the equilibrium measure for E.

Proof. First, recall that ν and μ_n for $n \geq 2$ are all supported on ∂E.

Now, let μ be any weak*-limit of the sequence of measures (μ_n). Then by the monotone convergence theorem, we have

$$I(\mu) = \lim_{m\to\infty} \int_{\partial E}\int_{\partial E} \max(\log|z-w|, -m) d\mu(z) d\mu(w).$$

Now, by the Stone-Weierstrass theorem, we have that $\mu_n \times \mu_n \to \mu \times \mu$ weak* on $\partial E \times \partial E$, so the above equality becomes

$$\begin{aligned} I(\mu) &= \lim_{m\to\infty}\lim_{n\to\infty} \int_{\partial E}\int_{\partial E} \max(\log|z-w|, -m) d\mu_n(z) d\mu_n(w) \\ &= \lim_{m\to\infty}\lim_{n\to\infty} \frac{1}{n^2}\sum_{j=1}^{n}\sum_{k=1}^{n} \max(\log|w_j^n - w_k^n|, -m) \\ &\geq \lim_{m\to\infty}\lim_{n\to\infty} \left(\frac{-m}{n} + \frac{2}{n^2}\sum_{j<k} \log|w_j^n - w_k^n|\right) \\ &= \lim_{n\to\infty} \frac{n-1}{n} \log \delta_n(E). \end{aligned}$$

This last limit is equal to $\log \mathrm{cap}(E) = I(\nu)$, by Theorem 2.7. The maximality and uniqueness of the equilibrium measure ν then implies that $\mu = \nu$, as required. □

Corollary 2.13. *For each $n \geq 2$, let $w_1^n, w_2^n, \ldots, w_n^n$ be a Fekete n-tuple for E. Then for each $z \in \Omega$,*

$$\frac{1}{n}\sum_{j=1}^{n} \log|z - w_j^n| \to p_\nu(z)$$

as $n \to \infty$.

In fact, the following result due to Pritsker yields uniform convergence on compact subsets of Ω as well as a precise rate of convergence, with the additional assumption that E is uniformly perfect.

Theorem 2.14 (Pritsker)**.** *Let E, $w_1^n, w_2^n, \ldots, w_n^n$, $n \geq 2$, be as above, and assume in addition that E is uniformly perfect. Then*

$$\left|\frac{1}{n}\sum_{j=1}^{n} \log|z - w_j^n| - p_\nu(z)\right| \leq C\frac{\log n}{\sqrt{n}} \qquad \left(z \in \Omega,\, g_\Omega(z,\infty) > \frac{1}{n}\right) \tag{2}$$

where $C = C(E)$ is a constant depending only on E.

Definition 2.15. We define the *Pritsker constant* of E to be the smallest constant $C(E)$ for which (2) holds.

We shall also need the following result, also due to Pritsker, on the rate of convergence of the n-th diameter to the logarithmic capacity.

Theorem 2.16 (Pritsker)**.** *Under the assumptions of Theorem 2.14, we have*

$$\log \frac{\delta_n(E)}{\operatorname{cap}(E)} \leq C_2 \frac{\log n}{\sqrt{n}},$$

where $C_2 = C_2(E)$ is a constant depending only on E.

See [14, Theorem 2.2] and its corollary.

3. Constructive approximation by Julia sets

In this section, we prove an explicit version of Theorem 1.2. As a consequence, we obtain Theorem 1.5 on the rate of approximation by Julia sets.

As before, let $E \subset \mathbb{C}$ be a compact set with connected complement Ω. We will construct a sequence of polynomials whose filled Julia sets totally approximate E, in the sense of Definition 1.3.

First, we can assume that the interior of E is not empty, adjoining a small disk in E if necessary. We can further suppose without loss of generality that 0 is an interior point of E, since otherwise it suffices to conjugate the polynomial by a suitable translation. Note that the above ensures that $\operatorname{cap}(E) > 0$, so that the Green's function $g_\Omega(\cdot, \infty)$ is well-defined.

We now introduce two positive geometric quantities associated with the set E, which we call the *inner radius* and *outer radius* of E. These are defined by

$$r(E) := \operatorname{dist}(0, \partial E)$$

and

$$R(E) := \sup_{w \in E} |w|$$

respectively. Note that $\overline{\mathbb{D}}(0, r(E)) \subset E \subset \overline{\mathbb{D}}(0, R(E))$.

Finally, let us assume as well that E is uniformly perfect. Recall that in this case, the Pritsker constant of E is defined to be the smallest constant $C(E)$ such that

$$\left| \frac{1}{n} \sum_{j=1}^{n} \log |z - w_j^n| - p_\nu(z) \right| \leq C(E) \frac{\log n}{\sqrt{n}} \qquad \left(z \in \Omega,\, g_\Omega(z, \infty) > \frac{1}{n} \right)$$

where $w_1^n, w_2^n, \ldots, w_n^n$ are the points of a Fekete n-tuple for E and ν is the equilibrium measure for E.

Theorem 3.1. *Let $s > 0$, and suppose that n is sufficiently large so that the following conditions hold :*

(i) $\dfrac{1}{n} \leq s$

(ii) $C(E) \dfrac{\log n}{\sqrt{n}} \leq \dfrac{s}{4}$

(iii) $e^{ns/4} \geq \dfrac{R(E) e^s}{r(E)}$

(iv) $\left(\dfrac{R(E) e^s}{r(E)} \right)^{1/n} \dfrac{\delta_n(E)}{\operatorname{cap}(E)} \leq e^{s/2}.$

Then the polynomial

$$P_n(z) = P_{n,s}(z) := z\frac{e^{-ns/2}}{\mathrm{cap}(E)^n}\prod_{j=1}^{n}(z - w_j^n)$$

satisfies

$$E \subset \mathcal{K}(P_n) \subset E_s,$$

where $E_s := E \cup \{z \in \Omega : g_\Omega(z, \infty) \leq s\}$.

In particular, the set E is totally approximable by the filled Julia sets of the polynomials P_n's, since the compact sets E_s shrink to E as s decreases to 0.

Proof. Let $\Omega_s := \widehat{\mathbb{C}} \setminus E_s = \{z \in \Omega : g_\Omega(z, \infty) > s\}$. Note that

$$\widehat{\mathbb{C}} \setminus \overline{\mathbb{D}}(0, R(E)e^s) \subset \Omega_s,$$

by Corollary 2.11.

Now, let $z \in \Omega_s$. Then $g(z, \infty) > 1/n$ by (i), so that by Theorem 2.14,

$$\left|\frac{1}{n}\sum_{j=1}^{n}\log|z - w_j^n| - p_\nu(z)\right| \leq C(E)\frac{\log n}{\sqrt{n}} \leq \frac{s}{4},$$

where we used condition (ii). In view of the relationship between $g_\Omega(z, \infty)$ and $p_\nu(z)$, we get

$$g_\Omega(z, \infty) - \frac{s}{4} \leq \frac{1}{n}\sum_{j=1}^{n}\log|z - w_j^n| - I(\nu)$$

and thus

$$\exp\left(n\left(g_\Omega(z, \infty) - \frac{s}{2} - \frac{s}{4}\right)\right) \leq |Q_n(z)|,$$

where

$$Q_n(z) := \frac{e^{-ns/2}}{\mathrm{cap}(E)^n}\prod_{j=1}^{n}(z - w_j^n) = \frac{P_n(z)}{z}.$$

Since $g_\Omega(z, \infty) > s$, we obtain

$$\frac{R(E)e^s}{r(E)} \leq e^{ns/4} < |Q_n(z)|, \tag{3}$$

by (iii), and this holds for all $z \in \Omega_s$.

On the other hand, if $z \in E$, then by Theorem 2.9 and (iv),

$$|Q_n(z)| \leq e^{-ns/2}\left(\frac{\delta_n(E)}{\mathrm{cap}(E)}\right)^n \leq e^{-ns/2}e^{ns/2}\frac{r(E)}{R(E)e^s} = \frac{r(E)}{R(E)e^s}. \tag{4}$$

Now, for $z \in \Omega_s$, we have, by (3),

$$|P_n(z)| = |z||Q_n(z)| > r(E)\frac{R(E)e^s}{r(E)} = R(E)e^s$$

so that $P_n(z) \in \widehat{\mathbb{C}} \setminus \overline{\mathbb{D}}(0, R(E)e^s) \subset \Omega_s$. Moreover, again by (3),

$$|P_n(z)| = |z||Q_n(z)| > |z|\frac{R(E)e^s}{r(E)},$$

where

$$\frac{R(E)e^s}{r(E)} > 1.$$

It follows that the iterates of P_n converge to ∞ on Ω_s, so that $\Omega_s \subset \widehat{\mathbb{C}} \setminus \mathcal{K}(P_n)$ and thus $\mathcal{K}(P_n) \subset E_s$.

Finally, for $z \in E$, we have, by (4),

$$|P_n(z)| = |z||Q_n(z)| \leq R(E)e^s \frac{r(E)}{R(E)e^s} = r(E)$$

and thus $P_n(E) \subset \overline{\mathbb{D}}(0, r(E)) \subset E$. Clearly, this implies that $E \subset \mathcal{K}(P_n)$, which completes the proof of the theorem. □

We can now use Theorem 3.1 to obtain a precise estimate for the rate of approximation. Recall from the introduction that $s_n(E)$, $n \in \mathbb{N}$, is defined to be the infimum of $s > 0$ for which there exists a polynomial p_n of degree n such that

$$E \subset \mathcal{K}(p_n) \subset E_s,$$

where $E_s := E \cup \{z \in \Omega : g_\Omega(z, \infty) \leq s\}$.

Solving for s in Theorem 3.1, one easily obtains the existence of a constant $c' = c'(E)$ such that

$$s_n(E) \leq c' \left(\frac{\log n}{\sqrt{n}} + \log \frac{\delta_n(E)}{\operatorname{cap}(E)} \right) \qquad (n \geq 5).$$

It follows from this and Theorem 2.16 that there is a constant $c = c(E)$ depending only on E such that

$$s_n(E) \leq c \frac{\log n}{\sqrt{n}} \qquad (n \geq 5),$$

thereby proving Theorem 1.5.

4. Hilbert's Lemniscate Theorem

In this section, we give a proof of Theorem 1.2 using a classical result of Hilbert on the approximation of planar sets by polynomial lemniscates. We first present a proof of this latter result based on Fekete polynomials.

Theorem 4.1 (Hilbert's Lemniscate Theorem [6])**.** *Let $E \subset \mathbb{C}$ be a compact set with connected complement Ω, and let U be an open neighborhood of E. Then there exists a polynomial Q such that*

$$|Q(z)| \leq 1 \qquad (z \in E)$$

and

$$|Q(z)| > 1 \qquad (z \in \mathbb{C} \setminus U).$$

Proof. Again, we can assume that $\operatorname{cap}(E) > 0$. Define

$$A := \inf_{z \in \widehat{\mathbb{C}} \setminus U} g_\Omega(z, \infty)$$

and

$$B := \sup_{z \in \widehat{\mathbb{C}} \setminus U} \tau_\Omega(z, \infty),$$

so that $A > 0$ and $B < \infty$. By Theorem 2.10, if q is a Fekete polynomial for E of degree n, then

$$\left(\frac{|q(z)|}{\|q\|_E} \right)^{1/n} \geq e^A \left(\frac{\operatorname{cap}(E)}{\delta_n(E)} \right)^B \qquad (z \in \mathbb{C} \setminus U).$$

Here we used the fact that $\delta_n(E) \geq \mathrm{cap}(E)$ for all n, cf. Theorem 2.7. Finally, since $\delta_n(E) \to \mathrm{cap}(E)$ as $n \to \infty$, the right-hand side will exceed 1 for all sufficiently large n, and setting $Q := q/\|q\|_E$ then gives the desired polynomial Q.

□

We can now use Theorem 4.1 to give another constructive proof of Theorem 1.2.

Proof. Again, we can assume without loss of generality that 0 belongs to E and that it is an interior point.

Let U be a small open neighborhood of E, and let Q be the polynomial as in Hilbert's theorem 4.1. Dividing Q by a constant slightly larger than one if necessary, we can assume that $\|Q\|_E < 1$.

Let $R, r > 0$ such that $\mathbb{D}(0,r) \subset E$ and $U \subset \mathbb{D}(0,R)$. For $k \geq 1$, define the polynomial $P_k(z) := zQ^k(z)$. Here Q^k denotes the k-th power of the polynomial Q. Then for k sufficiently large, we have

$$P_k(E) \subset \mathbb{D}(0,r) \subset E$$

and

$$P_k\left(\widehat{\mathbb{C}} \setminus U\right) \subset \widehat{\mathbb{C}} \setminus \mathbb{D}(0,R) \subset \widehat{\mathbb{C}} \setminus U,$$

since $P_k \to 0$ uniformly on E and $P_k \to \infty$ uniformly on $\widehat{\mathbb{C}} \setminus U$.

This implies that $E \subset \mathcal{K}(P_k)$. Also, the inequality

$$|P_k(z)| \geq |z| \min_{\widehat{\mathbb{C}} \setminus U} |Q| \qquad (z \in \widehat{\mathbb{C}} \setminus U),$$

where $\min\limits_{\widehat{\mathbb{C}} \setminus U} |Q| > 1$, shows that the iterates of P_k tend to ∞ on $\widehat{\mathbb{C}} \setminus U$, so that $\widehat{\mathbb{C}} \setminus U \subset \widehat{\mathbb{C}} \setminus \mathcal{K}(P_k)$, i.e. $\mathcal{K}(P_k) \subset U$.

Since U can be made arbitrarily close to E, the result follows.

□

5. Totally approximable sets

In this section, we prove Theorem 1.4, which states that a nonempty proper subset S of the complex plane $\mathbb{C}$ is totally approximable by filled Julia sets of polynomials of degree at least 2 if and only if S is bounded and $\mathbb{C} \setminus \mathrm{int}(S)$ is connected.

First, we need an approximation result.

Proposition 5.1. *Let S be a nonempty proper subset of the complex plane $\mathbb{C}$. If S is bounded and $\mathbb{C} \setminus \mathrm{int}(S)$ is connected, then S is totally approximable by a collection $\{K^\epsilon\}$ of nonempty compact sets with connected complement.*

The proof of Proposition 5.1 uses the following classical result from plane topology, known as Janiszewski's Theorem.

Lemma 5.2. *Let A and B be closed sets in $\widehat{\mathbb{C}}$ such that $A \cap B$ is connected. If A and B both have connected complement, then $A \cup B$ also has connected complement.*

Proof. See [13, Theorem 1.9]. □

We can now proceed with the proof of Proposition 5.1. For a set A and $\epsilon > 0$, we denote by $U_\epsilon(A)$ the open ϵ-neighborhood of the set A.

Proof. For $\epsilon > 0$, let G^ϵ be a grid of squares in $\mathbb{C}$ of sidelength $\epsilon/2$. Let H^ϵ be the union of the closed squares in G^ϵ which intersect ∂S. Note that since S is bounded, the set H^ϵ consists of finitely many closed squares and is therefore compact. Now, let F^ϵ be defined by taking each square Q in H^ϵ and replacing it by a smaller square $\widetilde{Q}$ with the same center of sidelength $\epsilon/4$. Then it is easy to see that the complement of F^ϵ is connected.

Now, define a collection of compact sets $\{K^\epsilon\}$ by

$$K^\epsilon := E^\epsilon \cup F^\epsilon,$$

where $E^\epsilon := \mathbb{C} \setminus U_\epsilon(\mathbb{C} \setminus \text{int}(S))$. Note that the sets E^ϵ and F^ϵ are disjoint, since F^ϵ is contained in the $(\epsilon/\sqrt{2})$-neighborhood of ∂S. See Figure 2.



FIGURE 2. On the left, a nonempty bounded subset S of the plane whose interior has connected complement. On the right, the corresponding approximating set K^ϵ.

We claim that for every $\epsilon > 0$, the complement of K^ϵ is connected. Indeed, first note that the complement of E^ϵ is connected. To see this, note that

$$\mathbb{C} \setminus E^\epsilon = \bigcup_{z \notin \text{int}(S)} \mathbb{D}(z, \epsilon),$$

hence each component of $\mathbb{C} \setminus E^\epsilon$ contains a disk $\mathbb{D}(z, \epsilon)$ for some $z \notin \text{int}(S)$ and therefore must intersect $\mathbb{C} \setminus \text{int}(S)$. Since $\mathbb{C} \setminus \text{int}(S)$ is connected and $\mathbb{C} \setminus E^\epsilon \supset \mathbb{C} \setminus \text{int}(S)$, it follows that each component of $\mathbb{C} \setminus E^\epsilon$ has to contain $\mathbb{C} \setminus \text{int}(S)$. Thus there can only be one component; in other words, the set $\mathbb{C} \setminus E^\epsilon$ is connected. Finally, since F^ϵ also has connected complement and E^ϵ, F^ϵ are disjoint, we can apply Lemma 5.2 to deduce that K^ϵ also has connected complement.

Next, we claim that S is totally approximable by the sets K^ϵ. It suffices to prove that $d(\partial S, \partial K^\epsilon) \to 0$ and $d(S, K^\epsilon) \to 0$ as $\epsilon \to 0$.

First, assume for a contradiction that $d(\partial S, \partial K^{\epsilon_n}) > \delta$ for all n, for some $\delta > 0$ and some sequence ϵ_n strictly decreasing to 0. Then for each n, either $\partial S \nsubseteq U_\delta(\partial K^{\epsilon_n})$ or $\partial K^{\epsilon_n} \nsubseteq U_\delta(\partial S)$. However, note that if $z \in \partial S$, then z belongs to a square Q in H^ϵ, so that the distance between z and the boundary of the corresponding square $\widetilde{Q}$ in F^ϵ is less than $\epsilon/(4\sqrt{2})$, hence $\partial S \subset U_\epsilon(\partial K^\epsilon)$, for all $\epsilon > 0$. We can therefore assume that the second case, $\partial K^{\epsilon_n} \nsubseteq U_\delta(\partial S)$, holds for all n. Now, recall that F^ϵ is contained in the $(\epsilon/\sqrt{2})$-neighborhood of ∂S. It follows that for all sufficiently large n, we have $\partial E^{\epsilon_n} \nsubseteq U_\delta(\partial S)$. Then for each such n, there is a point $z_n \in \partial E^{\epsilon_n} \subset \text{int}(S)$ such that the disk $\mathbb{D}(z_n, \delta)$ does not intersect the boundary of S, hence must be contained in $\text{int}(S)$. Passing to a

subsequence if necessary, we can assume that $z_n \to z_0 \in \overline{S}$. It is easy to see that z_0 necessarily belongs to ∂S, since the sets E^{ϵ_n} form a compact exhaustion of $\mathrm{int}(S)$ with $E^{\epsilon_n} \subset \mathrm{int}(E^{\epsilon_{n+1}})$ for all n. This contradicts the fact that for sufficiently large n, the disk $\mathbb{D}(z_0, \delta/2)$ is contained in the disk $\mathbb{D}(z_n, \delta)$, a subset of $\mathrm{int}(S)$. Therefore $d(\partial S, \partial K^\epsilon) \to 0$ as $\epsilon \to 0$.

It remains to prove that $d(S, K^\epsilon) \to 0$ as $\epsilon \to 0$. Again, assume for a contradiction that $d(S, K^{\epsilon_n}) > \delta$ for all n, for some $\delta > 0$ and some sequence ϵ_n strictly decreasing to 0. Since $E^\epsilon \subset \mathrm{int}(S)$ and F^ϵ is contained in the $(\epsilon/\sqrt{2})$-neighborhood of ∂S, we have $K^\epsilon \subset U_\epsilon(S)$, for all $\epsilon > 0$. We can therefore assume that for each n, $S \nsubseteq U_\delta(K^{\epsilon_n})$. Then for each n, there exists $z_n \in S$ such that the disk $\mathbb{D}(z_n, \delta)$ does not intersect K^{ϵ_n}. Assume without loss of generality that $z_n \to z_0 \in \overline{S}$. Then for all sufficiently large n, the disk $\mathbb{D}(z_0, \delta/2)$ does not intersect K^{ϵ_n}. If $z_0 \in \partial S$, then for sufficiently large n, the disk $\mathbb{D}(z_0, \delta/2)$ contains a square in F^{ϵ_n}, and we get a contradiction. If $z_0 \in \mathrm{int}(S)$, we get a contradiction with the fact that the sets E^{ϵ_n} form a compact exhaustion of $\mathrm{int}(S)$. This completes the proof of the proposition. □

Now, recall from Theorem 1.2 that any nonempty compact set with connected complement is totally approximable by polynomial filled Julia sets. Combining this with Proposition 5.1, we get that any nonempty bounded set whose interior has connected complement is also totally approximable by polynomial filled Julia sets. This proves the converse implication in Theorem 1.4.

For the proof of the other implication, we need the following elementary lemmas.

Lemma 5.3. *For any set $A \subset \mathbb{C}$, we have*

$$U_\epsilon(A) = A \cup U_\epsilon(\partial A).$$

Proof. Clearly, we have $A \subset U_\epsilon(A)$. Also, if $z \in U_\epsilon(\partial A)$, then $\mathbb{D}(z, \epsilon) \cap \partial A \neq \emptyset$, so that $\mathbb{D}(z, \epsilon) \cap A \neq \emptyset$, from which we deduce that $z \in U_\epsilon(A)$. This shows that $A \cup U_\epsilon(\partial A) \subset U_\epsilon(A)$.

For the other inclusion, if $z \in U_\epsilon(A)$ but $z \notin A$, then the disk $\mathbb{D}(z, \epsilon)$ intersects both A and $\mathbb{C} \setminus A$. By connectedness, it must intersect ∂A, so that $z \in U_\epsilon(\partial A)$. □

Lemma 5.4. *Let A, B be two nonempty proper subsets of the plane with $d(A, B) < \epsilon$ and $d(\partial A, \partial B) < \epsilon$. Then $d(\mathbb{C} \setminus A, \mathbb{C} \setminus B) \leq 2\epsilon$.*

Proof. We first show that $\mathbb{C} \setminus A \subset U_{2\epsilon}(\mathbb{C} \setminus B)$.

Let $z \in \mathbb{C} \setminus A$. If $z \notin U_\epsilon(A)$, then $z \in \mathbb{C} \setminus B$, since $B \subset U_\epsilon(A)$. On the other hand, if $z \in U_\epsilon(A)$, then by Lemma 5.3, we have that $z \in U_\epsilon(\partial A) \subset U_{2\epsilon}(\partial B)$. It follows that $\mathbb{D}(z, 2\epsilon) \cap \partial B \neq \emptyset$, so that $\mathbb{D}(z, 2\epsilon) \cap (\mathbb{C} \setminus B) \neq \emptyset$, from which we deduce that $z \in U_{2\epsilon}(\mathbb{C} \setminus B)$. In both cases, we get that $z \in U_{2\epsilon}(\mathbb{C} \setminus B)$.

This proves that $\mathbb{C} \setminus A \subset U_{2\epsilon}(\mathbb{C} \setminus B)$. The same argument with A and B interchanged yields the inclusion $\mathbb{C} \setminus B \subset U_{2\epsilon}(\mathbb{C} \setminus A)$, hence $d(\mathbb{C} \setminus A, \mathbb{C} \setminus B) \leq 2\epsilon$, as required.

□

We can now proceed with the proof of Theorem 1.4.

Proof of Theorem 1.4. Let S be a nonempty proper subset of $\mathbb{C}$.

As previously mentioned, if S is bounded and $\mathbb{C} \setminus \mathrm{int}(S)$ is connected, then by Proposition 5.1 and Theorem 1.2, the set S is totally approximable by filled Julia sets of polynomials of degree at least 2.

Conversely, assume that S is totally approximable by filled Julia sets of polynomials of degree at least 2. Then for any $\epsilon > 0$, there exists a polynomial P_ϵ such that $d(S, \mathcal{K}(P_\epsilon)) < \epsilon$ and $d(\partial S, \mathcal{J}(P_\epsilon)) < \epsilon$. Clearly, this implies that S is bounded, since for each $\epsilon > 0$, the set $\mathcal{K}(P_\epsilon)$ is compact.

It remains to prove that $\mathbb{C} \setminus \mathrm{int}(S)$ is connected. Assume for a contradiction that $\mathbb{C} \setminus \mathrm{int}(S) = E \cup F$, where E and F are disjoint nonempty closed subsets of $\mathbb{C}$. Since S is bounded, one of the sets E, F must be unbounded, say E, while the other set, F, is bounded. Now, fix a point $z_0 \in F$, and let $\delta > 0$ sufficiently small so that $\overline{\mathbb{D}}(z_0, 2\delta) \subset \mathbb{C} \setminus E$. Since $F \cup \overline{\mathbb{D}}(z_0, 2\delta)$ is a compact subset of the open set $\mathbb{C} \setminus E$, there is a bounded open set V with $F \cup \overline{\mathbb{D}}(z_0, 2\delta) \subset V \subset \overline{V} \subset \mathbb{C} \setminus E$. In particular, the boundary of V is contained in $\mathbb{C} \setminus (E \cup F) = \mathrm{int}(S)$.

Now, let $\epsilon > 0$ be smaller than both δ and half the distance between ∂V and $E \cup F$. Since $z_0 \in F \subset \mathbb{C} \setminus \mathrm{int}(S)$, we have that either $z_0 \in \partial S$ or $z_0 \in \mathbb{C} \setminus S$. In both cases, since $\partial S \subset U_\epsilon(\mathcal{J}(P_\epsilon))$ and $\mathbb{C} \setminus S \subset U_{2\epsilon}(\mathbb{C} \setminus \mathcal{K}(P_\epsilon))$ by Lemma 5.4, we get that the disk $\mathbb{D}(z_0, 2\epsilon)$ contains a point $w_0 \in \mathbb{C} \setminus \mathcal{K}(P_\epsilon)$.

This is enough to obtain a contradiction. Indeed, since $w_0 \in \mathbb{D}(z_0, 2\delta) \subset V$ and $w_0 \in \mathbb{C} \setminus \mathcal{K}(P_\epsilon)$, the connectedness of $\mathbb{C} \setminus \mathcal{K}(P_\epsilon)$ (see [11, Lemma 9.4]) implies that ∂V must have non-empty intersection with $\mathbb{C} \setminus \mathcal{K}(P_\epsilon)$. Since $\mathbb{C} \setminus \mathcal{K}(P_\epsilon) \subset U_{2\epsilon}(\mathbb{C} \setminus S)$, again by Lemma 5.4, it follows that the distance between ∂V and $\mathbb{C} \setminus \mathrm{int}(S) = E \cup F$ is less than 2ϵ, a contradiction.

Therefore, $\mathbb{C} \setminus \mathrm{int}(S)$ is connected, as required.

□

6. Numerical examples

In this section, we illustrate the method of Section 3 with some numerical examples.

As shown in Theorem 3.1, the computation of Julia sets approximating a given compact set E requires a numerical method for the computation of Fekete points for E. Unfortunately, the numerical computation of high-degree Fekete points is a hard large-scale optimization problem. Indeed, as far as we know, even for simple compact sets such as triangles, Fekete points have been computed only up to relatively small degrees, see e.g. [18], [12] and the references therein. In order to circumvent this issue, alternative numerical methods have been introduced for the computation of approximate Fekete points: see e.g. [17] for a method based on QR factorization of Vandermonde matrices with column pivoting. See also [3] for a discussion of the theoretical aspects of this method.

However, for slightly complicated sets, we were not able to use the algorithm of [17] to compute Fekete points of sufficiently high degree, due to rank-deficiency and ill-conditioning issues. For this reason, we decided to replace Fekete points by other equidistributed points, Leja points, which are much more convenient from a computational point of view. This sequence of points, introduced by Leja [8], is defined inductively as follows.

Choose an arbitrary point $z_1 \in \partial E$ and for $n \geq 2$, choose $z_n \in \partial E$ by maximizing the product

$$\prod_{j=1}^{n-1} |z - z_j|$$

for $z \in E$, i.e.

$$\prod_{j=1}^{n-1} |z_n - z_j| = \max_{z \in E} \prod_{j=1}^{n-1} |z - z_j|.$$

It is well-known that Leja points are easier to compute numerically than Fekete points, mainly because they are defined inductively by a univariate optimization problem. Furthermore, this optimization problem can be made discrete by replacing the compact set E by a sufficiently refined discretization.

We shall prove below that our approximation theorem 3.1 remains valid if Fekete points are replaced by Leja points, with small modifications. First, we need some lemmas.

As previously mentioned, Leja points are also equidistributed with respect to harmonic measure on ∂E, in the sense of Proposition 2.12. In fact, Pritsker's Theorem 2.14 also holds for Leja points, with the same constant.

Lemma 6.1. *If (z_n) is a sequence of Leja points for E, then*

$$\left| \frac{1}{n} \sum_{j=1}^{n} \log |z - z_j| - p_\nu(z) \right| \leq C(E) \frac{\log n}{\sqrt{n}} \qquad \left(z \in \Omega,\, g_\Omega(z, \infty) > \frac{1}{n} \right)$$

where $C(E)$ is Pristker's constant as in Definition 2.15.

Proof. See [14, Corollary 2.3]. □

We will also need an analogue of Theorem 2.7 for Leja points.

Lemma 6.2 (Leja)**.** *For $n \in \mathbb{N}$, define*

$$a_n := \prod_{j=1}^{n} |z_{n+1} - z_j|$$

Then $a_n^{1/n} \geq \operatorname{cap}(E)$ for each $n \in \mathbb{N}$, and $a_n^{1/n} \to \operatorname{cap}(E)$ as $n \to \infty$.

Proof. See [8, Lemme 1]. □

We can now prove the following variant of Theorem 3.1 for Leja points. As before, we assume that $E \subset \mathbb{C}$ is a uniformly perfect compact set with connected complement Ω and that $0 \in E$ is an interior point. Recall then that $r(E)$ and $R(E)$ are defined by

$$r(E) := \operatorname{dist}(0, \partial E)$$

and

$$R(E) := \sup_{w \in E} |w|.$$

Theorem 6.3 (Leja points version)**.** *Let $s > 0$, and suppose that n is sufficiently large so that the following conditions hold :*

(i) $\dfrac{1}{n} \le s$

(ii) $C(E)\dfrac{\log n}{\sqrt{n}} \le \dfrac{s}{4}$

(iii) $e^{ns/4} \ge \dfrac{R(E)e^s}{r(E)}$

(iv) $\left(\dfrac{R(E)e^s}{r(E)}\right)^{1/n} \dfrac{a_n^{1/n}}{\mathrm{cap}(E)} \le e^{s/2}$.

Then the polynomial

$$\widetilde{P}_n(z) = \widetilde{P}_{n,s}(z) := z\frac{e^{-ns/2}}{\mathrm{cap}(E)^n}\prod_{j=1}^{n}(z - z_j)$$

satisfies

$$E \subset \mathcal{K}(\widetilde{P}_n) \subset E_s,$$

where $E_s := E \cup \{z \in \Omega : g_\Omega(z, \infty) \le s\}$.

Remark. The only difference between the assumptions of Theorem 6.3 and those of Theorem 3.1 is condition (iv), where $\delta_n(E)$ is replaced by $a_n^{1/n}$.

Proof. The proof is exactly the same as in Theorem 3.1 except that in inequality (4), the use of Theorem 2.9 is replaced by the inequality

$$\prod_{j=1}^{n}|z - z_j| \le \prod_{j=1}^{n}|z_{n+1} - z_j| = a_n \qquad (z \in E),$$

which follows directly from the definition of Leja points.

□

The main improvement here is that the polynomials $\widetilde{P}_n$ are easier to compute numerically, even for high values of n. Indeed, the computation of $\widetilde{P}_n$ only relies on the computation of the first n Leja points for E, since the term $\mathrm{cap}(E)$ can be approximated by the quantity $a_n^{1/n}$, in view of Lemma 6.2. Now, given the polynomials $\widetilde{P}_n$, the computation of the corresponding Julia sets is quite standard and is easily done with MATLAB, for instance. We illustrate this with three examples.

Example 6.1.

In this example, the compact set E is the union of three components in the shape of a fish, a heart and a diamond.

Figure 3 is a representation of the filled Julia set of the polynomial

$$P_n(z) = z\frac{e^{-ns/2}}{\mathrm{cap}(E)^n}\prod_{j=1}^{n}(z - z_j),$$

with $n = 300$ and $s = 1/300$. Leja points were computed using a 30000 point discretization of E.



FIGURE 3. A filled Julia set in the shape of a fish, a heart and a diamond.

Example 6.2. In this example, the compact set E is the union of two components in the shape of the initials KL.

Figure 4 is a representation of the filled Julia set of the polynomial

$$P_n(z) = z\frac{e^{-ns/2}}{\mathrm{cap}(E)^n}\prod_{j=1}^{n}(z - z_j),$$

with $n = 300$ and $s = 1/300$. Leja points were computed using a 17000 point discretization of E.



FIGURE 4. A filled Julia set in the shape of the initials KL

Example 6.3. In this example, the compact set E is the union of two components in the shape of the initials MY.

Figure 5 is a representation of the filled Julia set of the polynomial

$$P_n(z) = z\frac{e^{-ns/2}}{\mathrm{cap}(E)^n}\prod_{j=1}^{n}(z - z_j),$$

with $n = 300$ and $s = 1/300$. Leja points were computed using a 22000 point discretization of E.

FIGURE 5. A filled Julia set in the shape of the initials MY

REFERENCES

1. V. Andrievskii, On the Approximation of a Continuum by Lemniscates, *J. Approx. Theory.*, **105** (2000), 292–304.
2. C.J. Bishop and K.M. Pilgrim, Dynamical dessins are dense, *Rev. Mat. Iberoam.*, **31** (2015), 1033–1040.
3. L. Bos and N. Levenberg, On the calculation of approximate Fekete points: the univariate case, *Electron. Trans. Numer. Anal.*, **30** (2008), 377–397.
4. L. Bos, N. Levenberg and S. Waldron, On the spacing of Fekete points for a sphere, ball or simplex, *Indag. Math. (N.S.)*, **19**(2008), 163–176.
5. M. Fekete, Über den transfiniten Durchmesser ebener Punktrnengen, *Math. Z.*, **32** (1930), 108–114.
6. D. Hilbert, Über die Entwicklung einer beliebigen analytischen Funktion einer Variabeln in eine unendliche nach ganzen rationalen Funktionen fortschreitende Reihe, *Göttinger Nachrichten*, (1897), 63–70.
7. Z. Laiyi, On the degree of convergence of lemniscates in finite connected domains, *J. Approx. Theory*, **131** (2004), 185–195.

8. F. Leja, Sur certaines suites liées aux ensembles plans er leur application à la représentation conforme, *Ann. Polon. Math.* **4** (1957), 8–13.
9. T. Kim, Quaternion julia set shape optimization, *Computer Graphics Forum*, **34** (2015), 157–176.
10. K.A. Lindsey, Shapes of polynomial Julia sets, *Ergodic Theory Dynam. Systems*, **35** (2015), 1913–1924.
11. J. Milnor, *Dynamics in one complex variable*, Princeton University Press, Princeton, 2006.
12. R. Pasquetti and F. Rapetti, Spectral element methods on unstructured meshes: Comparisons and recent advances, *J. Sci. Comput.*, **27** (2006), 377–387.
13. C. Pommerenke, *Univalent functions*, Vandenhoeck & Ruprecht, Göttingen, 1975.
14. I.E. Pritsker, Equidistribution of points via energy, *Ark. Mat.*, **47** (2011), 149–173.
15. T. Ransford, *Potential theory in the complex plane*, Cambridge University Press, Cambridge, 1995.
16. E. Saff and V. Totik, *Logarithmic potentials with external fields*, Springer-Verlag, Berlin, 1997.
17. A. Sommariva and M. Vianello, Computing approximate Fekete points by QR factorizations of Vandermonde matrices, *Comput. Math. Appl.*, **57** (2009), 1324–1336.
18. M.A. Taylor, B.A. Wingate, R.E. Vincent, An algorithm for computing Fekete points in the triangle, *SIAM J. Numer. Anal.*, **38** (2000), 1707–1720.

Department of Mathematics, University of Chicago, Chicago, IL 60637, United States.
E-mail address: klindsey@math.uchicago.edu

Department of Mathematics, Stony Brook University, Stony Brook, NY 11794-3651, United States.
E-mail address: malik.younsi@gmail.com